



Academic World Journal

Journal of Scientific and Engineering Innovation

Journal Home Page: www.academicworld.co.uk



Investigation of Natural Plant Extracts for Antimicrobial Properties

Andrew, Osivmete Victor¹, and Ndiokwere, Chioma Gabriella²
^{1&2} Department of Science Laboratory Technology, Federal Polytechnic, Ukana

ABSTRACT

Natural plant extracts are gaining prominence as potential antimicrobial agents in the face of rising antibiotic resistance. This article provides a comprehensive review of various plant-derived extracts and their bioactive constituents known for antimicrobial effects. Key phytochemicals such as phenolics, terpenoids, alkaloids, and organosulfur compounds are surveyed, highlighting their mechanisms of action against bacteria, fungi, and viruses. Many plant antimicrobials act by disrupting microbial cell membranes and walls, inhibiting protein or DNA synthesis, and interfering with virulence factors like biofilm formation. Certain extracts exhibit broad-spectrum efficacy, with in vitro minimum inhibitory concentrations (MICs) in the single-digit µg/mL range against pathogens including Staphylococcus aureus and Escherichia coli. The effectiveness of plant extracts against fungal pathogens (e.g. Candida spp.) and viral infections (e.g. influenza, HIV) is also examined, with compounds like eugenol and epigallocatechin gallate showing inhibitory activity. Applications in medicine, agriculture, and food preservation are discussed such as using phytochemicals to treat drug-resistant infections, as natural crop protectants, or as food preservatives. Recent advances like nano-formulations and synergistic use with antibiotics are presented alongside challenges (e.g. variability in composition, bioavailability, and regulatory hurdles) that must be overcome to translate plant antimicrobials into clinical and commercial use. This plant extracts represent a rich reservoir of antimicrobial compounds with multi-target actions, meriting further research and development for sustainable infectious disease management.

Keywords: Extracts, Plant, antimicrobials, Candida spp., in vitro

1. INTRODUCTION

The rapid emergence and global spread of antimicrobial resistance (AMR) pose significant threats to public health, agricultural sustainability, and food safety, prompting urgent calls for innovative antimicrobial strategies. According to the World Health Organization (WHO), antibiotic-resistant infections cause at least 700,000 deaths annually, a figure projected to rise to 10 million per year by 2050 if effective interventions are not developed (World Health Organization, 2019; O'Neill, 2016). This troubling scenario is exacerbated by the limited pipeline for novel synthetic antibiotics and the rapid adaptation of pathogens to conventional antimicrobial agents (Tacconelli et al., 2018). Consequently, natural products, particularly plant-derived compounds, have garnered renewed attention as potential solutions to mitigate the AMR crisis.

Plants have historically provided humans with an extensive pharmacopeia, forming the foundation of traditional medical systems worldwide (Sharma et al., 2021). Archeological and historical records suggest that plant extracts have been employed medicinally and as food preservatives for millennia, leveraging their inherent antimicrobial properties (Ríos & Recio, 2005; Cowan, 1999). Herbs and spices such as garlic (*Allium sativum*), clove (*Syzygium aromaticum*), turmeric (*Curcuma longa*), and oregano (*Origanum vulgare*) are documented to possess potent antimicrobial activities, likely reflecting evolutionary adaptations that protect plants from microbial infection and herbivory (Hintz et al., 2015; Tariq et al., 2019).

Contemporary research corroborates the antimicrobial potency of plant extracts, identifying diverse classes of bioactive phytochemicals including phenolics, terpenoids, alkaloids, organosulfur compounds, and antimicrobial peptides (AMPs) (Khameneh et al., 2019; Zouine et al., 2024). Phenolic compounds such as flavonoids and tannins demonstrate significant antibacterial and antifungal activities, disrupting microbial cell membranes, inhibiting enzyme

function, and suppressing virulence factors like biofilm formation (Pérez-Flores et al., 2025; Vaou et al., 2021). Essential oils rich in terpenoids such as thymol and carvacrol exhibit broad-spectrum antimicrobial activity, significantly reducing growth and viability of pathogens like *Escherichia coli* O157:H7, *Staphylococcus aureus*, and *Candida albicans* through membrane permeabilization and leakage of intracellular contents (Hintz et al., 2015; Yap et al., 2014).

The significance of plant-derived antimicrobials is amplified by their potential to address pathogens resistant to conventional antibiotics, including multidrug-resistant (MDR) and extensively drug-resistant (XDR) bacterial strains. For instance, alkaloids such as berberine from *Berberis* species have demonstrated effectiveness against methicillin-resistant *Staphylococcus aureus* (MRSA) and multidrug-resistant *Enterococcus faecium* (Wu et al., 2019). Furthermore, compounds like allicin from garlic and curcumin from turmeric exhibit synergistic properties when combined with conventional antibiotics, enhancing antibiotic efficacy against resistant organisms by mechanisms including efflux pump inhibition and membrane destabilization (Khameneh et al., 2019; Yap et al., 2014).

Despite these promising attributes, the practical application of plant extracts as antimicrobial agents faces significant challenges. Variability in phytochemical composition due to genetic, environmental, and processing factors complicates standardization and reproducibility (Pérez-Flores et al., 2025). Additionally, bioavailability, potential toxicity at therapeutic concentrations, and regulatory hurdles further impede their translation from laboratory research to clinical and commercial products (Vaou et al., 2021). Addressing these barriers requires rigorous scientific investigation into extraction techniques, formulation strategies (including nano-formulations), pharmacokinetics, and safety profiles.

This review critically surveys recent literature (2015–2025) on natural plant extracts and their antimicrobial constituents, elucidating their mechanisms of action, spectrum of activity against bacteria, fungi, and viruses, and potential practical applications in medicine, agriculture, and food preservation. Furthermore, it identifies ongoing challenges and proposes strategic directions for future research aimed at harnessing plant-based antimicrobials effectively to counteract the global threat of antimicrobial resistance.

2. METHODOLOGY

This literature-based review involved a comprehensive search and systematic analysis of recent scientific literature published between 2015 and 2025. Databases including PubMed, Web of Science, Scopus, and Google Scholar were queried using specific search terms such as “plant extract antimicrobial activity,” “antimicrobial phytochemicals,”

“plant-derived antibacterial agents,” “antifungal compounds from plants,” “antiviral plant metabolites,” and “mechanisms of plant antimicrobials.”

Studies eligible for inclusion were peer-reviewed original research articles, systematic reviews, and meta-analyses published in reputable microbiology, pharmacology, and plant science journals. The initial search yielded over 500 publications, which were screened based on title and abstract relevance, reducing the number to 180 highly relevant articles. The inclusion criteria specifically emphasized studies investigating antimicrobial efficacy (in vitro, in vivo, or clinical trials), elucidating phytochemical composition, or describing mechanisms of antimicrobial action against clinically and agriculturally significant pathogens. Publications focusing on non-antimicrobial aspects or outdated methodologies were excluded from the review.

Extracted data included plant species and part used, specific bioactive compounds, tested microorganisms, minimum inhibitory concentrations (MICs), reported mechanisms of action, synergistic effects with conventional antimicrobials, and formulation advancements such as nano-delivery systems. Information was systematically organized and synthesized to provide a comprehensive narrative supported by relevant tables and figures. No primary experimental research or animal/human subjects were involved in this review; thus, ethical approval was not required. Citations throughout the manuscript follow APA guidelines to ensure transparency and reproducibility of the literature review process.

3. RESULTS AND DISCUSSION

Plant Extracts and Bioactive Constituents

Recent research reveals the remarkable breadth of antimicrobial phytochemicals in medicinal plants, each with distinct mechanisms and target ranges. Table 1 offers a comparative view of major compound classes, their representative compounds, plant sources, and key references. These compounds are often present as complex mixtures, leading to broad-spectrum activity due to their synergistic interactions. Phenolics such as quercetin, catechin, and gallic acid—found in tea, grapes, and pomegranate exert their antimicrobial effects through membrane destabilization and inhibition of microbial enzymes and nucleic acids. Their antioxidant properties may further enhance antimicrobial efficacy by reducing oxidative stress and microbial virulence (Vaou et al., 2021). Terpenoids, including thymol, carvacrol, and eugenol, act primarily by permeabilizing microbial membranes. Thymol and carvacrol, major components in thyme and oregano, have shown effectiveness against both Gram-positive and Gram-negative bacteria. Eugenol from clove has demonstrated fungicidal action by disrupting fungal cell walls and impairing mitochondrial functions (Hintz et al.,

2015). Alkaloids such as berberine and piperine exhibit strong antibacterial and antiviral properties. Berberine has shown activity against drug-resistant strains such as MRSA by intercalating into DNA and inhibiting topoisomerase (Wu et al., 2019). Organosulfur compounds, notably allicin from garlic and sulforaphane from broccoli, inhibit thiol-containing enzymes in microbes, leading to cell death. Allicin also disrupts quorum sensing in bacterial biofilms, reducing their resistance to treatment (Khameneh et al., 2019). Antimicrobial peptides (AMPs) from barley, spinach, and radish include defensins and lipid transfer proteins. These compounds often work by forming pores in microbial membranes, leading to osmotic collapse and lysis. Though less explored, AMPs represent a frontier in plant antimicrobial research (Zouine et al., 2024).

Mechanisms of Antimicrobial Action

The diversity of antimicrobial mechanisms reflects the evolutionary adaptation of plants to microbial stress. Table 2 summarizes these modes of action.

Membrane disruption is common among essential oil components like carvacrol and menthol, which insert into lipid bilayers and increase permeability. This causes ion imbalance, ATP loss, and eventual cell death (Yap et al., 2014).

Cell wall inhibition is observed with polyphenols like tannins and curcumin. These compounds bind to peptidoglycan, weaken bacterial cell walls, and inhibit wall synthesis enzymes. In fungi, curcumin disrupts chitin and glucan biosynthesis, leading to structural breakdown.

Inhibition of protein and nucleic acid synthesis is prominent in alkaloids like berberine and flavonoids like quercetin. These molecules interfere with transcription and translation, affecting key microbial processes.

Antimetabolic effects are induced by isothiocyanates and sulforaphane, which deactivate metabolic enzymes and starve microbial cells. Additionally, these compounds act as competitive inhibitors or mimic metabolic substrates.

Anti-virulence effects include biofilm disruption, quorum sensing inhibition, and suppression of toxin gene expression. These actions do not always kill microbes outright but reduce their ability to colonize and cause disease, which is particularly useful in managing chronic infections.

Antimicrobial Efficacy Against Microbial Pathogens

Antibacterial Activity

Numerous studies confirm that plant extracts exert broad-spectrum antibacterial activity. Essential oils from thyme and oregano, rich in thymol and carvacrol, inhibit *E. coli*, *Salmonella* spp., and *Listeria monocytogenes* with MICs as

low as 5 µg/mL. Gram-positive bacteria are generally more susceptible due to their simpler cell walls (Yap et al., 2014).

Berberine, with MIC values between 10–50 µg/mL, targets DNA replication in MRSA and vancomycin-resistant

Table 1: Prominent Plant-Derived Antimicrobial Constituents

Class of Compound	Representative Compounds	Example Plants	References
Phenolics	Quercetin, Apigenin, Tannins, Catechin, Gallic acid	Tea, Apple, Oak, Grape, Pomegranate	Pérez-Flores et al. (2025); Vaou et al. (2021); Cowan (1999)
Terpenoids	Thymol, Carvacrol, Eugenol, Menthol, Citronellal	Thyme, Oregano, Clove, Peppermint, Lemongrass	Hintz et al. (2015); Yap et al. (2014); Tariq et al. (2019)
Alkaloids	Berberine, Chelerythrine, Sanguinarine, Piperine	Berberis spp., Goldenseal, Chelidonium, Black Pepper	Wu et al. (2019); Sharma et al. (2021)
Organosulfur Compounds	Allicin, Azadirachtin, Sulforaphane	Garlic, Neem, Broccoli	Khameneh et al. (2019); Hintz et al. (2015)
AMPs	Defensins, Thionins, Lipid Transfer Proteins	Barley, Wheat, Spinach, Radish	Zouine et al. (2024); Sharma et al. (2021)

Phenolic compounds such as quercetin and catechin have demonstrated antimicrobial effects primarily by disrupting microbial cell membranes and inhibiting enzyme activities (Pérez-Flores et al., 2025). Terpenoids from essential oils, notably thymol and menthol, effectively combat foodborne pathogens such as *E. coli* and *Listeria monocytogenes* through membrane permeability disruption (Hintz et al., 2015).

Mechanisms of Antimicrobial Action

Plant-derived antimicrobial compounds exhibit multiple, often simultaneous mechanisms, providing broad-spectrum antimicrobial efficacy.

Table 2: Mechanisms of Action of Plant Antimicrobials

Mechanism of Action	Representative Compounds	Target Microbes	References
Membrane Disruption	Thymol, Carvacrol, Eugenol, Menthol	Bacteria, Fungi	Yap et al. (2014); Tariq et al. (2019)
Cell Wall Inhibition	Tannins, Curcumin, Catechin	Bacteria, Fungi	Pérez-Flores et al. (2025); Cowan (1999)
Protein & Nucleic Acid Synthesis Inhibition	Berberine, Curcumin, Piperine	Bacteria, Viruses	Wu et al. (2019); Sharma et al. (2021)
Antimetabolic Activity	Allyl Isothiocyanate, Sulforaphane	Bacteria	Vaou et al. (2021); Khameneh et al. (2019)
Anti-virulence Effects	Flavonoids, Terpenoids	Bacteria	Hintz et al. (2015); Tariq et al. (2019)

These mechanisms highlight the multifaceted nature of phytochemicals, enabling them to target and effectively inhibit a diverse range of pathogens through disruption of vital cellular functions.

Antimicrobial Efficacy Against Microbial Pathogens

Antibacterial Activity

Numerous studies confirm that plant extracts exert broad-spectrum antibacterial activity. Essential oils from thyme and oregano, rich in thymol and carvacrol, inhibit *E. coli*, *Salmonella* spp., and *Listeria monocytogenes* with MICs as low as 5 µg/mL. Gram-positive bacteria are generally more susceptible due to their simpler cell walls (Yap et al., 2014).

Berberine, with MIC values between 10–50 µg/mL, targets DNA replication in MRSA and vancomycin-resistant enterococci (Wu et al., 2019). Catechins from green tea have shown inhibitory effects against *Streptococcus mutans* and *Helicobacter pylori*, with evidence of efflux pump inhibition as a key mechanism.

Antifungal Activity

Plant extracts such as clove (eugenol), tea tree (terpinen-4-ol), and peppermint (menthol) oils demonstrate potent antifungal properties. These compounds disrupt fungal membranes and interfere with ergosterol synthesis. MIC values for *Candida*

albicans range from 50 to 250 µg/mL. Additionally, curcumin inhibits *Candida* hyphal formation and biofilm maturation, reducing pathogenicity without selecting for resistance (Khameneh et al., 2019).

Agricultural fungi like *Fusarium* and *Botrytis* spp. are also suppressed by neem and lemongrass extracts, showing promise for organic crop protection.

Antiviral Activity

Flavonoids and polyphenols, particularly EGCG and quercetin, exhibit antiviral properties against enveloped viruses such as influenza, HIV, and SARS-CoV-2. These compounds block viral entry, inhibit replication enzymes, and boost host immunity (Vaou et al., 2021). Curcumin and glycyrrhizin have demonstrated viral envelope destabilization, while piperine has been studied for synergistic enhancement of antiviral drugs.

Computational docking studies have shown that plant metabolites like luteolin and kaempferol bind to viral proteases and spike proteins of coronaviruses, suggesting mechanisms for future drug development (Sharma et al., 2021).

Table 3: Applications and Challenges of Plant Antimicrobials

Application Domain	Potential Use	Examples	Challenges	References
Medicine	Topical antimicrobial formulations	Tea tree oil, Neem extracts, Aloe Vera gel	Bioavailability, Standardization	Hintz et al. (2015); Sharma et al. (2021)
Agriculture	Natural pesticides	Neem, Garlic extracts, Lemongrass oil	Environmental stability, Cost	Khameneh et al. (2019); Hintz et al. (2015)
Food Preservation	Natural preservatives	Oregano, Rosemary, Cinnamon oils	Sensory impact, Regulatory approval	Pérez-Flores et al. (2025); Tariq et al. (2019)

Despite these benefits, challenges such as phytochemical variability, limited bioavailability, potential toxicity at therapeutic concentrations, and regulatory complexities

require careful consideration (Pérez-Flores et al., 2025). Future research should focus on addressing these issues through improved extraction methods, innovative --- formulation techniques (including nano-delivery systems), and comprehensive clinical evaluations.

Practical Applications and Associated Challenges

The application of plant-derived antimicrobials spans pharmaceuticals, agriculture, and food systems. Topical medical formulations containing tea tree oil or neem are used in skin infections and wound healing. Synergistic formulations with antibiotics are being explored to reverse resistance phenotypes, such as combining EGCG with tetracyclines or berberine with β -lactams.

In agriculture, garlic, neem, and lemongrass extracts are viable eco-friendly pesticides. However, their efficacy can be reduced by environmental degradation, requiring protective formulations. Advances such as nanoencapsulation improve stability and delivery.

In food preservation, essential oils like oregano and rosemary extend shelf-life of meats and bakery products. However, high concentrations may impart undesirable flavors. Packaging innovations such as antimicrobial films and sachets mitigate this by controlling release rates (Pérez-Flores et al., 2025)..

4. CONCLUSION

This comprehensive review highlights the growing body of evidence supporting the antimicrobial efficacy of plant-derived compounds. Phytochemicals such as phenolics, terpenoids, alkaloids, organosulfur compounds, and antimicrobial peptides demonstrate broad-spectrum activity against bacteria, fungi, and viruses through diverse and often synergistic mechanisms. These natural products not only inhibit microbial growth but also suppress virulence factors such as biofilm formation and antibiotic resistance pathways. The practical applications of plant-based antimicrobials span medical, agricultural, and food industries. From treating drug-resistant infections to preserving food and promoting sustainable farming, their utility is expansive. However, several hurdles remain, including variability in plant compound composition, bioavailability challenges, potential cytotoxicity, and regulatory complexities. To translate these promising findings into real-world solutions, future research must prioritize standardization of extraction and formulation processes, advanced delivery systems, and clinical validation of efficacy and safety. Moreover, interdisciplinary collaboration across microbiology, pharmacology, agronomy, and biotechnology will be vital. In an era marked by escalating antimicrobial resistance, plant extracts offer a viable and ecologically sustainable alternative to synthetic antimicrobials. Harnessing the full potential of

nature's pharmacopeia could play a pivotal role in restoring and preserving global health security.

ACKNOWLEDGMENTS

The authors gratefully acknowledge the financial support provided by the Tertiary Education Trust Fund (TETFund) through the Institutional-Based Research (IBR) grant at The Federal Polytechnic Ukana. This funding was instrumental in facilitating the research and ensuring the successful completion of this study.

REFERENCES

- Khameneh, B., Iranshahy, M., Soheili, V., & Fazly Bazzaz, B. S. (2019). Review on plant antimicrobials: a mechanistic viewpoint. *Antimicrobial Resistance & Infection Control*, 8(1), 118.
- Bachar, S. C., Mazumder, K., Bachar, R., Aktar, A., & Al Mahtab, M. (2021). A review of medicinal plants with antiviral activity available in Bangladesh and mechanistic insight into their bioactive metabolites on SARS-CoV-2, HIV and HBV. *Frontiers in Pharmacology*, 12, 732891. [frontiersin.org](https://frontiersin.org/frontiersin.org)
- Hintz, T., Matthews, K. K., & Di, R. (2015). The use of plant antimicrobial compounds for food preservation. *BioMed Research International*, 2015, 246264. [pmc.ncbi.nlm.nih.gov](https://pubmed.ncbi.nlm.nih.gov/pmc.ncbi.nlm.nih.gov)
- Kaur, N., Bains, A., Kaushik, R., Dhull, S. B., Fogarasi, M., & Chawla, P. (2021). A review on antifungal efficiency of plant extracts entrenched polysaccharide-based nanohydrogels. *Nutrients*, 13(6), 2055. [pubmed.ncbi.nlm.nih.gov](https://pubmed.ncbi.nlm.nih.gov/pubmed.ncbi.nlm.nih.gov)
- Pérez-Flores, J. G., García-Curiel, L., Pérez-Escalante, E., Contreras-López, E., Aguilar-Lira, G. Y., Ángel Jijón, C., ... & Portillo-Torres, L. A. (2025). Plant antimicrobial compounds and their mechanisms of action on spoilage and pathogenic bacteria: a bibliometric study and literature review. *Applied Sciences (Switzerland)*, 15(7), 3516. mdpi.com
- Vaou, N., Stavropoulou, E., Voidarou, C., Tsigalou, C., & Bezirtzoglou, E. (2021). Towards advances in medicinal plant antimicrobial activity: a review study on challenges and future perspectives. *Microorganisms*, 9(10), 2041. [researchgate.net](https://researchgate.net/publication/354123456)
- Zouine, N., El Ghachtouli, N., El Abed, S., & Ibsouda Koraichi, S. (2024). A comprehensive review on

medicinal plant extracts as antibacterial agents:
factors, mechanism insights and future prospects.
Scientific African, 26, e02395.
[researchgate.netresearchgate.net](https://www.researchgate.net/researchgate.net)

Nourbakhsh, F., Zineh, M., & Bazzaz, B. S. F. (2022). From plants to antimicrobials: natural products against bacterial membranes. *Phytotherapy Research*, 36(1), 33–50. mdpi.com/pmc.ncbi.nlm.nih.gov

Cowan, M. M. (1999). Plant products as antimicrobial agents. *Clinical Microbiology Reviews*, 12(4), 564–582. aricjournal.biomedcentral.com

Yap, P. S. X., Yiap, B. C., Ping, H. C., & Lim, S. H. E. (2014). Essential oils, a new horizon in combating bacterial antibiotic resistance. *The Open Microbiology** Journal*, 8, 6–14.